

vals, if with blue they were totally blue, and so of the other Colours. And when they were illuminated with any one Colour, the Squares of their Diameters measured between their most luminous parts, were in the arithmetical progression of the numbers 0, 1, 2, 3, 4, and the Squares of the Diameters of their dark intervals in the progression of the intermediate numbers $\frac{1}{2}$, $1\frac{1}{2}$, $2\frac{1}{2}$, $3\frac{1}{2}$. But if the Colour was varied they varied their magnitude. In the red they were largest, in the indico and violet least, and in the intermediate Colours yellow, green and blue; they were of several intermediate bignesses answering to the Colour, that is, greater in yellow than in green, and greater in green than in blue. And hence I knew that when the Speculum was illuminated with white Light, the red and yellow on the outside of the Rings were produced by the least refrangible rays, and the blue and violet by the most refrangible, and that the Colours of each Ring spread into the Colours of the neighbouring Rings on either side, after the manner explained in the first and second Part of this Book, and by mixing diluted one another so that they could not be distinguished, unless near the center where they were least mixed. For in this Observation I could see the Rings more distinctly, and to a greater number than before, being able in the yellow Light to number eight or nine of them, besides a faint shadow of a tenth. To satisfy my self how much the Colours of the several Rings spread into one another, I measured the Diameters of the second and third Rings, and found them when made by the confine of the red and orange to be the same Diameters when made by the confine of blue and indico, as 9 to 8, or thereabouts. For it was hard

to determine this
cles made success
differed more from
fively by the gree
made by the viole
on the computatio
differences of the
outmost red, the
of orange and yel
the confine of gre
indico, the confine
let, are in propor
of a Monochord
sol, la, fa, sol, la, m
 $\frac{1}{16}, \frac{1}{12}, \frac{1}{12}, \frac{2}{27}, \frac{1}{27}, \frac{1}{18}$ A
by the confine of
the Circle made b
8 A as above, the
the difference of t
the outmost red, a
as $\frac{1}{16} + \frac{1}{12} + \frac{1}{12} + \frac{2}{27}$ t
the difference of th
and by the confine
to $\frac{1}{12} + \frac{1}{18}$, that is, a
fore these differen
first to 9 A and f
will have the Di
least and most refi
Diameters are the
50 to 41, and the
as 3 to 2 very nea
much from the p